

Citation for published version:

Woodman, TJ, Wood, PJ, Thompson, AS, Hutchings, TJ, Jiao, P, Steel, GR, Threadgill, MD & Lloyd, MD 2012, 'Chiral inversion of NSAID-CoA esters by human -methylacyl-CoA racemase', Biochemical Society Conference, 2012, Cambridge, UK United Kingdom, 1/01/12.

Publication date:
2012

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Chiral inversion of NSAID-CoA esters by human α -methylacyl-CoA racemase 1A

T. J. Woodman, P. J. Wood, A. S. Thompson, T. Hutchings, P. Jiao, G. R. Steel, M. D. Threadgill, **M. D. Lloyd**

Medicinal Chemistry, Department of Pharmacy & Pharmacology, University of Bath, Claverton Down, Bath BA2 7AY, United Kingdom; M.D.Lloyd@bath.ac.uk

Ibuprofen and other 2-arylpropanoic acids (2-APAs) are non-steroidal anti-inflammatory drugs (NSAIDs) and the 2S- enantiomers act by inhibition of cyclo-oxygenase 1 and 2 (COX-1 and -2). 2-APA drugs are given as racemic mixtures and chiral inversion is essential for pharmacological activity. This pathway for 2*R*-ibuprofen consists of conversion to the 2*R*-acyl-CoA ester, chiral inversion to the 2*S*-epimer, and hydrolysis to 2*S*-ibuprofen. The chiral inversion reaction is catalysed by α -methylacyl-CoA racemase (AMACR; P504S). AMACR is increased ~10-fold in prostate and other cancers, and is a novel cancer drug target. Other 2-APAs undergo chiral inversion *in vivo*, but it is unclear if AMACR is involved.

The present study investigates whether AMACR catalyses chiral inversion of other 2-APA-CoA esters. Substrates were incubated with human AMACR 1A in $^2\text{H}_2\text{O}$ -containing buffer and all exchanged the 2-proton, an obligatory step for chiral inversion. Derivatization of products from chiral substrates showed a *ca.* 1:1 epimeric mixture of products, consistent with the proposed deprotonation/reprotonation mechanism. Steady state kinetic analysis showed that most substrates were converted with similar efficiency ($k_{\text{cat}}/K_{\text{m}} = 100\text{--}150 \text{ M}^{-1} \text{ s}^{-1}$). Fenoprofenoyl-CoA was converted much more efficiently ($k_{\text{cat}}/K_{\text{m}} = \sim 1426 \text{ M}^{-1} \text{ s}^{-1}$), due to its low K_{m} value ($\sim 2.3 \text{ }\mu\text{M}$ *cf.* 26–74 μM).

The results suggest that all 2-APA drugs undergo the same chiral inversion pathway. 2-APAs may be useful for treatment of cancers with increased AMACR levels.

222 words